



Your checklist in serious situations!

1. Treat

Treat the patient's head as quickly as possible with mosquito® med Head Lice Shampoo 10.

2. Family

All members of the family must be carefully checked for nits and head lice and should also be treated with mosquito® med Head Lice Shampoo 10 on a precautionary basis.

3. Laundry

In order to ensure that laundry is free from head lice, towels, hats, pillowcases and other linen must be washed at a minimum of 60 °C.

4. Cleaning

Combs and hairbrushes can be cleaned with a hand brush in a hot soapy solution. Other non-washable items that have come in contact with the patient's head (e.g., bicycle helmets) should be stored in a sealed plastic bag for three days - head lice cannot survive longer without a food source.

5. Notify

You must notify kindergartens, schools and all persons with whom your child comes in contact. This can help stop the further spread of head lice.

6. Check

Afterwards carry out daily checks. If you find any live head lice, you must immediately repeat the treatment with mosquito® med Head Lice Shampoo 10. Even if you do not find any more head lice - you must repeat the treatment after 7 to 10 days.

7. Protect

The mosquito® lice repellent spray can be used as a preventive measure. It prevents head lice head lice infestation for up to 12 hours.



Head Lice Shampoo 10
100 ml
PZN 10415469



Also available in a
200 ml family size bottle
PZN 10415475



Head Lice Repellent Spray
100 ml
PZN 10834982



Head lice comb
with integrated magnifying lens
PZN 01873902

1 Use mosquito® lice protectant with caution. Before use always read the label and product information carefully.



WEPA Apothekenbedarf GmbH & Co KG
D-56204 Hillscheid
www.wepa-dieapothekenmarke.de

www.mosquito-laese.de